

VARIATIONS IN HAEMATOLOGICAL PARAMETERS IN OBESE, SEDENTARY, AEROBIC-  
TRAINING AND RESISTANCE-TRAINING POPULATIONS

Betül Kaplan<sup>1</sup>, Hakim Çelik<sup>2</sup>, Yusuf Hoşoğlu<sup>3</sup>, Mehmet Küçük<sup>2</sup>, Yakup Aktaş<sup>4</sup>, Mehmet Ali Eren<sup>5</sup>

**ABSTRACT**

**Introduction:** Although physical activity has a considerable impact on various laboratory markers, evidence on haematological alterations, cardiovascular risk factors and inflammatory mediators after resistance-training is limited. **Objective:** The aim of this study is to investigate alterations in hematological parameters among populations engaging in obesity, sedentary behavior, aerobic exercise, and resistance training. **Materials and Methods:** A study with 122 male volunteers aged 18-45 divided into four groups based on exercise type and Body Mass Index. It compared blood parameters of resistance exercisers to normal-weight sedentary, overweight/obese individuals, and normal-weight aerobic exercisers. Blood count included RBC, HCT, reticulocyte, platelet counts, hemoglobin, MCV, MCH, RDW, and MPV. Neutrophil-to-lymphocyte ratio (NLR), platelet-to-lymphocyte ratio (PLR), and systemic immune-inflammation index (SII) were determined based on hemogram. **Results:** Resistance training may have negative effects on individuals at cardiovascular risk, but it was found to be advantageous for athletes or sedentary adults with a good vascular endothelium to raise HCT and, possibly, enhance tissue blood supply by inducing vasodilation.

**Key words:** Sedentary. Obese. Body mass index.

1 - Dialysis Program, Hasan Kalyoncu University, Gaziantep, Türkiye.

2 - Department of Physiology, Harran University, Sanliurfa, Türkiye.

3 - Education and Research Hospital, Adiyaman, Turkey.

4 - School of Physical Education and Sports, Harran University, Sanliurfa, Türkiye.

5 - Department of Endocrinology, Harran University, Sanliurfa, Türkiye.

**RESUMO**

**Variações nos parâmetros hematológicos em populações obesas, sedentárias, praticantes de treinamento aeróbico e de treinamento de resistência**

**Introdução:** Embora a atividade física tenha um impacto significativo em vários marcadores laboratoriais, as evidências sobre alterações hematológicas, fatores de risco cardiovascular e mediadores inflamatórios após o treinamento de resistência são limitadas. **Objetivo:** O objetivo deste estudo é investigar as alterações nos parâmetros hematológicos em populações envolvidas em obesidade, comportamento sedentário, exercício aeróbico e treinamento de resistência. **Materiais e Métodos:** Um estudo com 122 voluntários do sexo masculino, com idades entre 18 e 45 anos, divididos em quatro grupos com base no tipo de exercício e índice de massa corporal. Foram comparados os parâmetros sanguíneos de praticantes de exercícios de resistência com sedentários de peso normal, indivíduos com sobrepeso/obesidade e praticantes de exercícios aeróbicos de peso normal. A contagem sanguínea incluiu contagens de RBC, HCT, reticulócitos, plaquetas, hemoglobina, VCM, HCM, RDW e VPM. A razão neutrófilo-linfócito (NLR), a razão plaqueta-linfócito (PLR) e o índice imune-inflamatório sistêmico (SII) foram determinados com base no hemograma. **Resultados:** O treinamento de resistência pode ter efeitos negativos em indivíduos com risco cardiovascular, mas mostrou-se vantajoso para atletas ou adultos sedentários com um bom endotélio vascular, aumentando o HCT e, possivelmente, melhorando o suprimento sanguíneo tecidual por meio da indução de vasodilatação.

**Palavras-chave:** Sedentário. Obeso. Índice de massa corporal.

E-mail dos autores:

betul.tatlibem@hku.edu.tr

hakimcell@gmail.com

yhosoglu@gmail.com

mhmtkck81@gmail.com

drmalieren@harran.edu.tr

## INTRODUCTION

Physical activity is essential for the promotion of health. However, incorrect exercise could be hazardous, and cardiovascular complications during exercise could result in sudden death (Kimura et al., 2015).

Many epidemiologic studies have shown that exercise protects against coronary artery disease (CAD). A recent meta-analysis revealed that moderate to vigorous physical exercise is connected with a decreased risk of coronary artery disease. In particular, intense and moderate exercise reduced the incidence of coronary artery disease by 27% and 11%, respectively, as compared to persons with minimal or no physical activity (Sofi et al., 2008).

Several variables, such as dehydration, an increase in hematocrit, plasma viscosity, red blood cell (RBC) aggregation, and a decrease in RBC deformability caused by intense exercise, may contribute to a decrease in blood fluidity (Kimura et al., 2015).

Resistance exercise training (RET) combined with dietary protein supplementation is a frequent technique among athletes and leisure exercisers with the goal of improving RET-induced increases in muscle development and strength (Morton et al., 2018).

Muscle injury brought on by exercise triggers an augmented immune response, leading to the production of inflammatory signalling molecules from leukocytes and their various cellular subtypes. The final stage of this process is the entry of inflammatory cells into the injured region (Cardoso et al., 2012).

There is a correlation between high-intensity exercise and a biphasic shift in the number of circulating leukocytes. An increase in the overall number of leukocytes is seen in the immediate post-exercise phase. This rise in the total number of leukocytes happens mostly at the cost of lymphocytes, neutrophils, and, to a lesser degree, monocytes (Cardoso et al., 2012).

Atherosclerosis is linked to the presence of persistent inflammation at a low grade. Inflammatory cells, such as white blood cells (WBC), are critical to the progression of atherosclerosis in the arterial artery wall. This disease may be prevented by eliminating inflammatory cells. Because of this, having a high WBC count is linked to an increased risk of developing cardiovascular illnesses (Madjid, et al., 2004; Ross, 1999).

Obesity is a chronic metabolic disorder that is related with cardiovascular disease as well as increased morbidity and mortality rates (Coban et al., 2005). In recent years, it has come to be regarded as the nation's important public health problem and an epidemic (Wyrostek et al., 2021).

A numerous studies have shown that obesity and being overweight are connected with an increased platelet reactivity (Leite et al., 2016; Ranucci et al., 2019; Vilahur, Ben-Aicha, Badimon, 2017).

As a result, official guidelines propose that each person should undertake cardiorespiratory exercise for 20–60 minutes three–five days each week. These guidelines are based on research that shown that strenuous physical activity decreases the risk of cardiovascular disease (Lamprecht et al., 2013).

We planned a prospective research to examine the changes of some parameters of the complete blood cell count (CBC) in resistance exercisers, normal-weight sedentary, normal-weight regular exercisers, and overweight or obese in order to determine whether the increased cardiovascular problems and inflammation that are sometimes observed after certain kinds of exercise may be reflected by variation of red cell distribution width (RDW), mean platelet volume (MPV) and Systemic immune-inflammation index (SII).

There was no other research in the literature that compared the hematological effects of resistance exercise to those of obesity and regular exercise.

## MATERIALS AND METHODS

This research was conducted at the Harran University Faculty of Medicine, Department of Physiology laboratory, and 122 healthy adult male participated in this study. All individuals were informed of the objectives of the study and permission was acquired prior to their participation.

The Harran University Clinical Research Ethics Committee gave its authorization to carry out this research in a decision that was dated 05.09.2022 and numbered 2022/17/27.

The research was carried out in a manner that was compliant with both the Declaration of Helsinki and the standards for Good Clinical Practice. After receiving ethical approval and having a signed permission form

on file, we started recruiting participants. Authors had access to information that could identify individual participants during or after data collection.

Individuals were placed into one of four categories based on their preferred type of exercise and their body mass index (BMI);

**(1) Resistance Exercise Group (REG):** This group consisted of 31 healthy male volunteers that participate in bodybuilding on a regular basis and have a BMI that is greater than what is considered normal (28,11). The mean duration of exercise the participants was determined as  $6.45 \pm 5.45$  (year  $\pm$  standard deviation)

**(2) Sedentary Group (SG):** This group consisted of 29 healthy male volunteers who did not exercise regularly (Sedentary) and had a normal BMI (22.70).

**(3) Obesity and Overweight Group (OG):** This group consisted of 30 healthy male volunteers who did not participate in any regular activities and whose body mass index was much higher than the average (28.75).

**(4) Regular Aerobic Exercise Group (AEG):** This group consisted of 32 healthy male volunteers with a normal body mass index (22.56) who participated in regular aerobic exercise in the form of brisk walking and running on a regular basis. The mean duration of exercise the participants was determined as  $5.93 \pm 3.10$  (year  $\pm$  standard deviation).

#### Collection of Blood and Hemogram Measurement

Following the collection of the essential demographic data, 2 ml of venous blood was drawn from each volunteer participant following an 8-hour overnight fasting into tubes containing ethylenediaminetetraacetic acid. Within 2 hours of blood withdrawal, the Abbott Cell-DYN Ruby Hematology Analyzer (USA) was used to analyse whole blood. Red blood cell (RBC), haematocrit (HCT), reticulocyte, and platelet counts, haemoglobin, mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), RBC distribution width (RDW), and mean platelet volume (MPV) were included in the complete blood count. Also, systemic immune-inflammation index (SII), the neutrophil-to-lymphocyte ratio (NLR), the platelet-to-lymphocyte ratio (PLR), and the

HGB-to-RDW ratio were calculated. Total neutrophil and platelet counts were divided by lymphocyte counts to get the NLR and PLR, respectively. The peripheral platelet (P), neutrophil (N), and lymphocyte (L) counts were used to arrive at the Systemic immune-inflammation index (SII) using the formula  $SII = (P \cdot N)/L$  (Chen et al., 2017).

#### Statistical Analysis

For statistical analysis, the Windows-compatible IBM SPSS 25.0 (IBM SPSS Inc, Chicago, IL, USA) package program was used. Using the Shapiro-Wilk test, skewness, kurtosis, histogram, and Q-Q plot, we assessed the conformity of the data to the normal distribution. Continuous variables following the normal distribution were represented by the Mean (M)  $\pm$  Standard Deviation (SD), whereas continuous variables not following the normal distribution were expressed as the Median (interquartile range of values). Between groups, non-normally distributed data were analysed with the Kruskal Wallis H test (Bonferroni correction was used for within-group comparisons), and normally distributed data were analysed with one-way analysis of variance (Tukey's test for within-group comparisons). In the analysis, 95% was accepted as the confidence interval.  $p < 0.05$  was statistically significant.

#### RESULTS

There was no statistically significant difference in the height and age values of the groups when their demographic information was evaluated (Table 1).

As predicted, the BMI and weights of obese and muscle-obesity resistance exercisers were considerably greater than those of lean groups (CE and REG).

The results of the study's hemogram analysis are outlined in Table 2, which provides a summary of the findings. In the groups that were under investigation, there were no significant changes found in WBC, NEU, LYM, HGB, MCH, RDW and PLT. Similar outcomes were observed for inflammatory indexes such as SII, NLR, PLR, and the HGB-to-RDW ratio.

**Table 1 - Demographic data.**

	REG (n=31) Mean ± SD	SG (n=29) Mean ± SD	OG (n=30) Mean ± SD	AEG (n=32) Mean ± SD	p
Age (Year)	28,55 ± 4,83	27,10 ± 5,98	28,67 ± 5,49	26,63 ± 5,52	0.357
Height (cm)	176,45 ± 6,63	177,21 ± 5,28	178,00 ± 5,90	179,38 ± 5,68	0.244
Weights (kg)	87,62 ± 7,17 <sup>a,c</sup>	71,34 ± 7,40 <sup>d</sup>	91,22 ± 8,34 <sup>f</sup>	72,62 ± 6,11	<0.001
BMI (kg/m <sup>2</sup> )	28,11 ± 0,99 <sup>a,c</sup>	22,70 ± 1,97 <sup>d</sup>	28,75 ± 1,64 <sup>f</sup>	22,56 ± 1,43	<0.001

Variables were represented by the Mean (M) ± Standard Deviation (SD). RGE: Resistance Exercise Group, SG: Sedentary Group, OG: Obesity and Overweight Group, AEG: Regular Aerobic Exercise Group. a: Defined as a statistical difference between REG and SG. c: Defined as a statistical difference between REG and AEG. d: Defined as a statistical difference between SG and OG. f: Defined as a statistical difference between OG and AEG.

**Table 2 - Hematological Parameters.**

	REG (n=31) Mean ± SD	SG (n=29) Mean ± SD	OG (n=30) Mean ± SD	AEG (n=32) Mean ± SD	p
WBC	8,55 (3,00)	7,67 (2,29)	7,65 (2,64)	6,86 (2,02)	0,134
NEU	4,58 (1,91)	4,08 (1,71)	3,94 (1,53)	3,92 (1,12)	0,771
LYM	2,54 (1,73)	2,39 (0,58)	2,77 (1,06)	2,31 (0,77)	0,073
MONO	0,67 (0,20) <sup>c</sup>	0,64 (0,22)	0,54 (0,18)	0,51 (0,20)	0,005
HGB	16,60 (1,50)	16,50 (1,60)	16,30 (1,27)	16,40 (4,03)	0,618
HCT	53,15 ± 4,03 <sup>a,b,c</sup>	50,32 ± 3,12	50,74 ± 3,21	50,96 ± 2,50	0,005
MCV	87,25 (7,90) <sup>b</sup>	87,70 (3,79)	85,30 (3,80)	87,60 (5,28)	0,024
MCH	28,00 (1,70)	28,35 (1,80)	27,50 (2,17)	28,50 (2,10)	0,253
MCHC	31,80 (2,30) <sup>b,c</sup>	32,20 (0,90)	32,50 (1,45)	32,45 (1,83)	0,018
RDW	11,55 (1,93)	10,90 (1,30)	11,15 (1,03)	11,00 (0,78)	0,564
PLT	268,44 ± 50,06	283,40 ± 46,60	288,60 ± 50,25	277,53 ± 59,79	0,495
MPV	6,96 ± 1,38 <sup>a</sup>	7,84 ± 1,19	7,64 ± 1,27	7,39 ± 1,00	0,038
NLR	1,56 (0,90)	1,67 (0,63)	1,37 (0,56)	1,69 (0,58)	0,390
PLR	108,36 (61,82)	112,90 (36,30)	102,34 (49,02)	123,95 (29,04)	0,203
SII	426,51 (395,13)	501,19 (245,01)	418,57 (163,12)	482,71 (202,67)	0,602
HGB/RDWR	1,46 ± 0,17	1,46 ± 0,16	1,47 ± 0,13	1,48 ± 0,15	0,897

Variables were represented by the Mean (M) ± Standard Deviation (SD) or Median (interquartile range of values, IQR). RGE: Resistance Exercise Group, SG: Sedentary Group, OG: Obesity and Overweight Group, AEG: Regular Aerobic Exercise Group. a: Defined as a statistical difference between REG and SG. b: Defined as a statistical difference between REG and OG. c: Defined as a statistical difference between REG and AEG.

## DISCUSSION

There is compelling evidence that unhealthy lifestyle factors, such as smoking, alcohol consumption, physical inactivity, and being overweight or obese, contributed to approximately 60% of deaths and were linked to chronic inflammation status, highlighting the potential modifiability of the effects of chronic inflammation by these factors (Li et al., 2021).

It has been shown in recent years that the neutrophil-to-lymphocyte ratio (NLR), platelet-to-lymphocyte ratio (PLR), and systemic immune-inflammation index (SII) are all valid indicators of inflammatory responses at the cellular level.

Some studies have shown a rise in NLR in response to acute or intense exercise (Korkmaz et al., 2018; Schlagheck et al., 2020; Wahl et al., 2020).

However, prolonged (chronic) exercise regimens appear to have contradictory findings. In the study conducted by Makras et al., (2005) on healthy male adults, four weeks of military training consisting of intermittent moderate activity decreased neutrophils and the neutrophil-to-lymphocyte ratio.

In the study conducted by Wang, Chen, Chen (2011), on obese male teenagers, a 4-week diet and activity intervention dramatically decreased the neutrophil to lymphocyte ratio. The intervention was linked with a considerable reduction in pro-inflammatory cytokine concentrations.

In contrast, there was no significant difference between the obese, resistance exercise, sedentary, and regular exercise groups in this study. The absence of long-term change in blood neutrophil and lymphocyte levels may be related to the body's adaptability to this environment, despite the possibility of acute increases in blood neutrophil and lymphocyte levels in chronically applied regular exercise regimens.

PLR levels were significantly affected by high intensity exercise, with values almost twice as high as at rest. Possible explanations include exercise-dependent platelet mobilization in peripheral circulation (Walzik, Joisten, 2021). Regarding the effect of persistent exercise on the PLR, after 3 weeks of endurance exercise in a group with multiple sclerosis, no changes were seen (Joisten et al., 2021). Similarly, there was no significant difference in PLR values between the chronic

exercise groups and the obese group in our investigation.

Recently, the systemic immune-inflammation index (SII), which is derived using neutrophil, lymphocyte, and platelet counts in peripheral blood, was proposed as a measure of the equilibrium between systemic inflammation and immunological state. Similar to NLR, SII readings are highest when neutrophil and platelet numbers are high and lymphocyte numbers are low.

Prior research suggested that a high SII level was related with a worse outcome for individuals with certain cancers (Li et al., 2021).

Also, It was hypothesized that chronic inflammation, which might cause reactive nitrogen species, reactive oxygen species, genomic instability, and cell senescence, increased the risk of cardiovascular disease and death in participants with higher SII levels (Li et al., 2021).

Schlagheck et al., (2020) who aimed to evaluate the differences in cellular immunological alterations between acute endurance exercise and resistance exercise, determined that resistance exercises have no effect on SII.

Discovered no significant changes in the mean values of NLR, SII, and PLR between the overweight obese and severe obese weight groups (Țaranu et al., 2021).

In our study, resistance exercisers were compared to the obese, regular exercise, and sedentary groups, however there was no significant difference between the groups.

During exercise recovery, under the effect of glucocorticoids, substantial numbers of lymphocytes and monocytes enter the circulation. Monocytes activated by exercise are likely to penetrate skeletal muscle and develop into tissue-resident macrophages that enable repair and regeneration, especially following strenuous exercise sessions that produce severe skeletal muscle damage.

Additionally, monocytes with effector characteristics are redeployed preferentially after exercise (Peake, Neubauer, 2017). Among the interesting findings of our study is that the monocyte count was highest in the resistance-exercising group. This increase suggested that resistance workouts, which place a greater demand on the skeleton's muscular tissue, can increase the number of monocytes.

MPV, an indication of platelet activation, plays a significant role in the pathogenesis of cardiovascular disorders (Varol

et al., 2010). MPV is a crucial biological characteristic, and bigger platelets possess a greater thrombogenic potential (Martin, 1990). Based on the findings of this study, resistance exercise has been found to lower MPV and may lessen the thrombogenic potential of blood.

Several studies have shown that exercise increases haematocrit and blood viscosity. It is hypothesized that many mechanisms, including fluid shift, water loss, release of sequestered RBCs from the spleen, and water retention in the muscle, are responsible for this increase (Connes et al., 2013).

Young erythrocytes, the number of which rises as a consequence of the hemolysis that occurs during exercise, have a lower MCHC as well as a higher MCV and deformability.

Also, it is well established that a lower MCV and a higher MCHC both contribute to an increase in the internal viscosity of RBCs, which in turn reduces their deformability (Temiz et al., 2000).

Kilic-Toprak et al., (2012) reported an increase in haematocrit and blood viscosity following 12 weeks of progressive resistance training.

In a different study of cycling exercise, researchers discovered that cycling often raises blood viscosity by attributable to increases in plasma viscosity and haematocrit (Brun et al., 1993).

In contrast, aerobic exercise research demonstrates that extended outdoor walking does not affect blood viscosity or haematocrit (Neuhaus, Gaehtgens, 1994; Tripette et al., 2011).

High blood viscosity has generally been believed to have a detrimental effect on performance from an exercise physiology standpoint. An excessive rise in blood viscosity during exercise has been hypothesized to be potentially harmful to the cardiovascular system because it is believed to enhance vascular resistance and the post-load work of the heart (Connes et al., 2006; Connes et al., 2013; Yalcin et al., 2003).

But it is also known that the increased wall shear stress may induce nitric oxide (NO) generation by endothelial cells to provide a vasodilatory compensation in response to an increase in blood viscosity (Cabrales et al., 2006; Vázquez et al., 2011).

The study's limitations include the fact that it did not use a bigger sample size by

including other cardiovascular imaging modalities, blood viscosity, and other inflammation indicators.

These restrictions may be considered in future research. Nevertheless, this compensating reaction is contingent on the health condition of the endothelium, and a rise in blood viscosity may be more harmful to the vascular system in the event of endothelial dysfunction (Salazar Vázquez et al., 2010).

## CONCLUSION

According to the findings of our study, resistance training may have negative effects on individuals at cardiovascular risk, but it was found to be beneficial for athletes or sedentary individuals with a healthy vascular endothelium to increase HCT and thereby increase tissue blood supply by stimulating vasodilation. In addition, prolonged resistance workouts did not produce any rise in the inflammatory indices assessed in the blood in our study, and it was determined to be crucial for the body's defence since it increases the number of monocytes, which then spread to the tissues as macrophages.

## REFERENCES

- 1-Brun, J.; Fons, C.; Supparo, I.; Mallard, C.; Orsetti, A. Could exercise-induced increase in blood viscosity at high shear rate be entirely explained by hematocrit and plasma viscosity changes? *Clinical Hemorheology and Microcirculation*. Vol. 13. Num. 2. 1993. p, 187-199.
- 2-Cabrales, P.; Martini, J.; Intaglietta, M.; Tsai, A.G. Blood viscosity maintains microvascular conditions during normovolemic anemia independent of blood oxygen-carrying capacity. *American Journal of Physiology-Heart and Circulatory Physiology*. Vol. 291. Num. 2. 2006. p. H581-H590.
- 3-Cardoso, A.; Bagatini, M.; Roth, M.; Martins, C.; Rezer, J.; Mello, F.; Schetinger, M. Acute effects of resistance exercise and intermittent intense aerobic exercise on blood cell count and oxidative stress in trained middle-aged women. *Brazilian journal of medical and biological research*. Num. 45. 2012. p. 1172-1182.

- 4-Chen, J.H.; Zhai, E.T.; Yuan, Y.J.; Wu, K.M.; Xu, J.B.; Peng, J.J.; Cai, S.R. Systemic immune-inflammation index for predicting prognosis of colorectal cancer. *World J Gastroenterol*. Vol. 23. Num. 34. 2017. p. 6261-6272. doi: 10.3748/wjg.v23.i34.6261
- 5-Coban, E.; Ozdogan, M.; Yazicioglu, G.; Akcıt, F. The mean platelet volume in patients with obesity. *International journal of clinical practice*. Vol. 59. Num. 8. 2005. p. 981-982.
- 6-Connes, P.; Sara, F.; Hardy-Dessources, M.-D.; Marlin, L.; Etienne, F.; Larifla, L.; Hue, O. Effects of short supramaximal exercise on hemorheology in sickle cell trait carriers. *European journal of applied physiology*. Vol. 97. Num. 2. 2006. p. 143-150.
- 7-Connes, P.; Simmonds, M.J.; Brun, J.-F.; Baskurt, O.K. Exercise hemorheology: classical data, recent findings and unresolved issues. *Clinical Hemorheology and Microcirculation*. Vol. 53. Num. 1-2. 2013. p. 187-199.
- 8-Joisten, N.; Proschinger, S.; Rademacher, A.; Schenk, A.; Bloch, W.; Warnke, C.; Zimmer, P. High-intensity interval training reduces neutrophil-to-lymphocyte ratio in persons with multiple sclerosis during inpatient rehabilitation. *Multiple Sclerosis Journal*. Vol. 27. Num. 7. 2021. p. 1136-1139.
- 9-Kilic-Toprak, E.; Ardic, F.; Erken, G.; Unver-9-Kocak, F.; Kucukatay, V.; Bor-Kucukatay, M. Hemorheological responses to progressive resistance exercise training in healthy young males. *Med Sci Monit*. Vol. 18. Num. 6. 2012. p. Cr351-360. doi: 10.12659/msm.882878
- 10-Kimura, T.; Hamada, H.; Taito, S.; Takahashi, M.; Sekikawa, K. The effect of exercise on blood fluidity: use of the capillary model to assess the cloginess of blood. *Clinical Hemorheology and Microcirculation*. Vol. 61. Num. 4. 2015. p. 559-569.
- 11-Korkmaz, A.; Yıldız, A.; Türker Duyuler, P.; Duyuler, S.; Yılmaz, S.; Basyigit, F.; Ileri, M. Combination of change in hematological parameters with exercise stress test to predict coronary artery disease. *Journal of clinical laboratory analysis*. Vol. 32. Num. 1. 2018. p. e22205.
- 12-Lamprecht, M.; Moussalli, H.; Ledinski, G.; Leschnik, B.; Schlagenhauf, A.; Koestenberger, M.; Cvirn, G. Effects of a single bout of walking exercise on blood coagulation parameters in obese women. *Journal of applied physiology*. Vol. 115. Num. 1. 2013. p. 57-63.
- 13-Leite, N.R.P.; Siqueira de Medeiros, M.; Mury, W.V.; Matsuura, C.; Perszel, M.B.M.; Noronha Filho, G.; Mendes-Ribeiro, A.C. Platelet hyperaggregability in obesity: is there a role for nitric oxide impairment and oxidative stress? *Clinical and Experimental Pharmacology and Physiology*. Vol. 43. Num. 8. 2016. p. 738-744.
- 14-Li, H.; Wu, X.; Bai, Y.; Wei, W.; Li, G.; Fu, M.; Guo, H. Physical activity attenuates the associations of systemic immune-inflammation index with total and cause-specific mortality among middle-aged and older populations. *Sci Rep*. Vol. 11. Num. 1. 2021. p. 12532. doi: 10.1038/s41598-021-91324-x
- 15-Madjid, M.; Awan, I.; Willerson, J.T.; Casscells, S.W. Leukocyte count and coronary heart disease: implications for risk assessment. *J Am Coll Cardiol*. Vol. 44. Num. 10. 2004. p. 1945-1956. doi: 10.1016/j.jacc.2004.07.056
- 16-Makras, P.; Koukoulis, G.N.; Bourikas, G.; Papatheodorou, G.; Bedevis, K.; Menounos, P.; Kartalis, G. Effect of 4 weeks of basic military training on peripheral blood leucocytes and urinary excretion of catecholamines and cortisol. *Journal of sports sciences*. Vol. 23. Num. 8. 2005. p. 825-834.
- 17-Martin, J. Platelet heterogeneity in vascular disease *Platelet Heterogeneity*. p. 205-226: Springer. 1990.
- 18-Morton, R.W.; Murphy, K.T.; McKellar, S.R.; Schoenfeld, B.J.; Henselmans, M.; Helms, E.; Phillips, S.M. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. Vol. 52. Num. 6. 2018. p. 376-384. doi: 10.1136/bjsports-2017-097608
- 19-Neuhaus, D.; Gaehtgens, P. Haemorheology and long term exercise. *Sports Medicine*. Vol. 18. Num. 1. 1994. p. 10-21.

- 20-Peake, J.M.; Neubauer, O. Recovery of the immune system after exercise. *Vol. 122. Num. 5. 2017. p. 1077-1087. doi: 10.1152/jappphysiol.00622.2016*
- 21-Ranucci, M.; Aloisio, T.; Di Dedda, U.; La Rovere, M.T.; Arroyabe, B.M.L.; Baryshnikova, E. Platelet reactivity in overweight and obese patients undergoing cardiac surgery. *Platelets. Vol. 30. Num. 5. 2019. p. 608-614.*
- 22-Ross, R. Atherosclerosis-an inflammatory disease. *N Engl J Med. Vol. 340. Num. 2. 1999. p. 115-126. doi: 10.1056/nejm199901143400207*
- 23-Salazar Vázquez, B.Y.; Salazar Vázquez, M.A.; Jáquez, M.G.; Bracho Huemoeller, A.H.; Intaglietta, M.; Cabrales, P. Blood pressure directly correlates with blood viscosity in diabetes type 1 children but not in normals. *Clinical Hemorheology and Microcirculation. Vol. 44. Num. 1. 2010. p. 55-61.*
- 24-Schlagheck, M.L.; Walzik, D.; Joisten, N.; Koliymitra, C.; Hardt, L.; Metcalfe, A.J.; Zimmer, P. Cellular immune response to acute exercise: Comparison of endurance and resistance exercise. *Eur J Haematol. Vol. 105. Num. 1. 2020. p. 75-84. doi: 10.1111/ejh.13412*
- 25-Sofi, F.; Capalbo, A.; Cesari, F.; Abbate, R.; Gensini, G.F. Physical activity during leisure time and primary prevention of coronary heart disease: an updated meta-analysis of cohort studies. *European Journal of Preventive Cardiology. Vol. 15. Num. 3. 2008. p. 247-257.*
- 26-Țaranu, I.; Lazea, C.; Creț, V.; Răcățianu, N.; Iancu, M. Inflammation-Related Markers and Thyroid Function Measures in Pediatric Patients: Is the Grade of Obesity Relevant?. *Vol. 11. Num. 3. 2021. doi: 10.3390/diagnostics11030485*
- 27-Temiz, A.; Başkurt, O.; Pekcetin, C.; Kandemir, F.; Güre, A. Leukocyte activation, oxidant stress and red blood cell properties after acute, exhausting exercise in rats. *Clinical Hemorheology and Microcirculation. Vol. 22. Num. 4. 2000. p. 253-259.*
- 28-Tripette, J.; Hardy-Dessources, M.-D.; Beltan, E.; Sanouiller, A.; Bangou, J.; Chalabi, T.; Nebor, D. Endurance running trial in tropical environment: A blood rheological study. *Clinical Hemorheology and Microcirculation. Vol. 47. Num. 4. 2011. p. 261-268.*
- 29-Varol, E.; Akcay, S.; Ozaydin, M.; Erdogan, D.; Dogan, A.; Altinbas, A. Mean platelet volume is associated with insulin resistance in non-obese, non-diabetic patients with coronary artery disease. *Journal of cardiology. Vol. 56. Num. 2. 2010. p. 154-158.*
- 30-Vázquez, B.Y.S.; Cabrales, P.; Tsai, A.G.; Intaglietta, M. Nonlinear cardiovascular regulation consequent to changes in blood viscosity. *Clinical Hemorheology and Microcirculation. Vol. 49. Num. 1-4. 2011. p. 29-36.*
- 31-Vilahur, G.; Ben-Aicha, S.; Badimon, L. New insights into the role of adipose tissue in thrombosis. *Cardiovascular research. Vol. 113. Num. 9. 2017. p. 1046-1054.*
- 32-Wahl, P.; Mathes, S.; Bloch, W.; Zimmer, P. Acute impact of recovery on the restoration of cellular immunological homeostasis. *International Journal of Sports Medicine. Vol. 41. Num. 01. 2020. p. 12-20.*
- 33-Walzik, D.; Joisten, N. Transferring clinically established immune inflammation markers into exercise physiology: focus on neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio and systemic immune-inflammation index. *Vol. 121. Num. 7. 2021. p. 1803-1814. doi: 10.1007/s00421-021-04668-7*
- 34-Wang, R.; Chen, P.; Chen, W. Diet and exercise improve neutrophil to lymphocyte ratio in overweight adolescents. *International Journal of Sports Medicine. Vol. 32. Num. 12. 2011. p. e1-e5.*
- 35-Wyrostek, J.; Piotrowska, A.; Czerwińska-Ledwig, O.; Zuziak, R.; Szyguła, Z.; Cisoń, T.; Pilch, W. Complex effects of whole body cryostimulation on hematological markers in patients with obesity. *PloS one. Vol. 16. Num. 4. 2021. p. e0249812.*
- 36-Yalcin, O.; Erman, A.; Muratli, S.; Bor-Kucukatay, M.; Baskurt, O.K. Time course of hemorheological alterations after heavy anaerobic exercise in untrained human subjects. *Journal of applied physiology. Vol. 94. Num. 3. 2003. p. 997-1002.*



**RBNE**  
**Revista Brasileira de Nutrição Esportiva**

---

Corresponding author  
Betül Kaplan  
betul.tatlibadem@hku.edu.tr  
Dialysis Program.  
Hasan Kalyoncu University Havaalanı Yolu  
Üzeri 8. km. Şahinbey / GAZİANTEP  
Zip code: 27000

Recebido para publicação em 07/03/2024

Aceito em 11/09/2024